



Global Advanced Research Journal of Medicinal Plant (GARJMP) Vol. 3(1) pp. 010-014, February 2015  
Available online <http://garj.org/garjmp/index.htm>  
Copyright © 2015 Global Advanced Research Journals

*Full Length Research Paper*

## Prevalance of tuberculosis in District Dir Upper, Pakistan

**Naveed Khan, Sardar Azhar Mehmood, Muzafar Shah\*, Shabir Ahmad, Sajid Mahmood and Mazhar Ali**

Department of zoology, Hazara University Mansehra-21300, Pakistan

Accepted 19 January, 2015

The present study was carried out to investigate the prevalence of tuberculosis (TB) in sub division Sheringal, Dir proper, Barawal and Wari District Dir Upper in Khyber Pakhtunkwa (KP) province, Pakistan from (January 2013 to December 2013). During this survey I collect data from four tehsils of District Dir Upper. A total of laboratory reports of 380 patients were collected from four different tehsils, in which 98 (25.78%) were found positive and 282 (74.21%) were found negative tuberculosis patients. The higher incidence of tuberculosis was showed in tehsil sheringal 31 (28.18%). And lowest rate of tuberculosis was found in tehsil wari 18 (22.5%). And in Gender wise prevalence out of 98 positive cases 37 (22.42%) were male and 67 (28.37%) were female. This shows that the rate of tuberculosis is higher in female as compare to male. The higher infection of tuberculosis were found in the people of middle age from 13-30 (70.58%) and lowest infection are found in the children and adult about (8.57%). In month wise the higher prevalence was recorded in the month of March (46.15%) while the lowest prevalence was found in the month of April showing (16.21%).

**Keywords:** Tuberculosis, Prevalence, Khyber Pakhtunkhwa.

### INTRODUCTION

Tuberculosis (TB) is a bacterial disease which is a major cause of morbidity and mortality throughout the world. It remained the world's leading cause of death since decades. TB becomes active disease and in 75% cases it affects lungs i.e. pulmonary tuberculosis while 25% cases suffer from extra-pulmonary tuberculosis. The main symptoms include three weeks prolonged and productive cough, chest pain and coughing up blood (WHO, 2006). Tuberculosis is caused by *Mycobacterium tuberculosis* which is slow growing facultative intracellular parasite.

*Mycobacterium tuberculosis* is being existed in human population since ancient times, first infected humans 10,000-15,000 years ago (Ducati *et. al.*, 2006). However it was in the seventeenth century that pathological and anatomical descriptions of tuberculosis (TB) disease began to appear (National Tuberculosis Center, 2009). An estimated one third of the world's population is infected with *Mycobacterium tuberculosis*, presenting a major impediment to tuberculosis control. When the World Health Organization declared TB a global health emergency in 1992, it was prevalent in almost all countries of the world. WHO estimated a total of 9.27 million new cases worldwide in 2007 with 1.3 million deaths with >90% in developing countries (WHO, 2008).

\*Corresponding Author E-mail: [muzafarphd@hotmail.com](mailto:muzafarphd@hotmail.com)

Although 95% of TB positive cases and 97% of all deaths occur in high-endemic countries, the disease continued to be a problem in industrialized countries as well, mostly in immigrant population, in elderly individuals with reactivating latent infection, and in local outbreaks (Broekmans *et. al.*, 2002).

## AIMS AND OBJECTIVES

The objective of my study is,

- To find out the prevalence of Tuberculosis in District Dir Upper.
- To study the prevalence of TB among individuals of different age groups.

## METHOD AND MATERIALS

### Study Area and Map

District Dir is located in Khyber Pukhtunkhwa, Pakistan. The total area is 5282 km<sup>2</sup>. It lies from 34° 38' north in latitude and from 71° 30' to 72° 22' east in longitudes. District Dir is divided into two Districts, Dir Upper and Dir Lower. I collect data from District Dir Upper from four different tehsils. Tehsil Sheringal, Tehsil Dir, Tehsil Brawl, and Tehsil Wari. Dir Upper is one of 24 districts of Pukhtunkhwa, which was founded in 1916. Entire District lies in valley of the Panjkora river which rises in the Hindokash. The population of the area is 1,375,657. The annual rainfall is 400 mm, which is recorded in summer (UNICEF, 2008).

### Material

Following material was used to conduct my research work.

Questionnaires, Sputum collection tubes, Slides, AFB test reagents, Microscope etc.

### Methodology

This investigation has been conducted among different Tehsils of District Upper Dir at DHQ Upper Dir. The group of patients were selected within the age range of 1-60 years and above, patients having clinical features suggestive of tuberculosis as evening pyrexia, weight loss, productive cough, haemoptysis, malaise, tiredness, anorexia, chest pain, and patients with raised ESR and chest X-ray finding abnormal shadows, cavitations, and/or abnormality in the lymph nodes relative symptoms to tuberculosis were included.

A printed Performa containing a comprehensive record of all patients was filled after getting necessary information from each patient. Blood with ESR, Urine R/E, Chest X-ray, Blood Urea, Blood Sugar, and AFB smear examination by direct microscopy were done for all patients.

AFB (Acid-Fast Bacilli) smear refers to the microscopic examination of a fluorochrome stain of a clinical specimen. The sputum specimen was placed on the slide with the help of a wire loop to make a smear with the help of forceps and then acid-fast stain was put on the sample followed by carbon fuchsin for 5 min to fix it on the slide. Then the slide was washed with decolorizer for about 3 minutes. Then methylene blue was added for a minute to examine under microscope using 100 X eye piece.

For more confirmation I also used ICT test which was carried out through blood serum collected after centrifugation of blood at high speed. 10 µL of serum was transferred to a special tube designed for ICT test followed by addition of 3 drops of buffer solution for 15 minutes. When the T line turned dark, it revealed the presence of TB infection.

### Statistical Analysis

The data was statistically analyzed and tabulated using Microsoft Excel.

## RESULTS

The present study was conducted on the prevalence of tuberculosis (TB) in District Dir Upper. This study was based on laboratory reports of patients from different hospitals. Total laboratory reports were collected and grouped into age wise, area wise, and gender wise. A total of laboratory reports of 380 patients were collected, in which 98 were found positive and 282 were found negative tuberculosis patients. Four Tehsils Sheringal, Dir, Brawl, and Wari of District Upper Dir were screened for prevalence of TB. The higher incidence of tuberculosis was shown in Tehsil Sheringal 31 (28.18%). And lowest rate of tuberculosis was found in Tehsil Wari 18 (22.5%) (Table 1, Figure 1). And in gender wise prevalence out of 98 positive cases 37 (22.42%) were male and 67 (28.37%) were female (Table 2, Figure 2). This shows that the rate of tuberculosis is higher in female as compared to male. In age wise the higher infection of tuberculosis were found in the people of middle age from 13-30 (70.58%) and lowest infection are found in the children and adult about (8.57%) (Table 3, Figure 3). In month wise the higher prevalence was recorded in the month of March (46.15%) while the lowest prevalence was found in the month of April showing (16.21%) (Table 4, Figure 4).

**Table 1.** Prevalence of tuberculosis in different tehsils of District Dir Upper, Pakistan from January to December 2013

Area	Total	Positive	%	Negative	%
Sheringal	110	31	28.18	79	71.81
Dir	100	25	25	75	75
Wari	80	18	22.5	62	77.5
Brawal	90	24	26.66	66	73.33
Total	380	98	25.78	282	74.21

**Table 2.** Prevalence of Tuberculosis patients in gender wise from January to December 2013

Gender	Total	Positive	Negative	%+ve	%-ve
Male	165	37	128	22.42	77.57
Female	215	61	154	28.37	71.62
Total	380	98	282	31.63	68.63

**Table 3.** Prevalence of tuberculosis in age wise of District Dir Upper, Pakistan from January to December 2013

Age	Total samples	Total +ive	%	-ive	%
1-12	70	6	8.57	64	91.42
13-30	85	60	70.58	25	29.41
31-40	80	13	16.25	67	83.75
41-50	75	6	8	69	92
51-60	40	8	20	32	80
Above 60	30	5	16.66	25	83.33
Total	380	98	25.78	282	74.21

**Table 4.** Prevalence of tuberculosis in month wise of District Dir Upper, Pakistan from January 2013 to December 2013.

Months	Total samples	Positive	%age	Negative	%age
January	41	10	24.39	31	75.60
February	31	8	25.80	23	74
March	26	12	46.15	14	53.84
April	37	6	16.21	31	83.78
May	33	8	24.24	25	75.75
June	22	7	31.81	15	68
July	25	5	20	20	80
August	33	8	24.24	25	75.75
September	31	6	19.35	25	80.64
October	26	8	30.76	18	69
November	33	8	24.24	25	75.75
December	42	12	28.57	30	71.42
Total	380	98	25.78	282	74.21

## DISCUSSION

It has been confirmed by present study that smear-positive tuberculosis has been found on individuals of low socioeconomic group and in females. The patient's family

is being always on the risk of transmit the tuberculosis infection. Tuberculosis has caused tremendous amount of worldwide than any other infectious disease especially in developing countries like Pakistan where it is declared fourth major cause of death. There were 181/100,000

estimated new cases and 223/100,000 prevalent cases in Pakistan according to 2007 estimates (Nisar, 2007). Early diagnosis and effective treatment of active cases that are infectious to the community is the best way of controlling TB in our country. The main reason of the increased risk of the manifestation of the disease could possibly be the delay in diagnosis and failure to cure a great percentage of smear positive cases, which might lead to the high death toll ratio and MDR cases in Pakistan. Currently for diagnosis, developing countries rely on AFB stains and culture and radiographic changes. In this study total of 380 patients were investigated for active TB.

Majority of the infected persons were from middle class. Common signs at presentation were anemia and had bronchial breathing. During investigation most of the patients especially female were having Hb level of 8–10 gm/dl. As they remain most functional reproductively in the age between 13 to 30 years their personal health may be compromised. The higher prevalence in female is also due to longer to seek care (patient delay) due to stigma and social exclusion, heavier workloads, prioritization of other family members over own well-being, lack of independence, inaccessibility to financial resources and powerlessness in decision making. They experience longer provider diagnostic and treatment delays. They are engaged in more activities that need to be replaced in the household. In addition, women have higher direct costs than men, because they often need somebody to accompany them; they are less mobile and have less financial resources.

Similarly, TB is highly prevalent in age groups ranging from 13 to 30 years, and this is due to their extensive exposure to the outside environment characterized by poor sanitation and unhygienic conditions. Individuals of this age group remain in frequent contact with industrial environments where they inhale air rich in effluents and dust particles which may cause alveolar inflammation resulting in immune suppression.

The conclusion of this study that lockup convicts are at amplified risk for TB infection is in line with previous studies driven in the parts of the world. It has been proposed that a TB control program should be introduced. There should be special attention for the personages showing positive family history for TB as they have already provoked a chance to gain the manifestation of the disease. Also, deprived sanitation and freshening provide *M. tuberculosis*, a grand opportunity to persist for extensive periods and transmit it to others. The improvement within ventilation, sanitation, and overall living conditions in our societies/houses environment have been suggested thoroughly. We endorse an awareness program for families to acquire knowledge about the possibility for the transmission of *M. tuberculosis* infection from AFB sputum smear-positive pulmonary tuberculosis.

## CONCLUSION AND RECOMMENDATIONS

The investigation report provides pulsating statistics on the TB infection condition amongst the District Dir Upper population. TB vestiges a foremost public health problem, focus the tribal population, and there is a need of further TB Control programs on a constant and study basis. Targeted programs must engage prisons, jails, HIV/STD programs, specific high-risk occupational groups, illicit drug users, and persons living on the street. Smear positive pulmonary TB is more prevalent in females, in young age individuals, and in people of low socio-economic group. For the control of tuberculosis, early diagnosis of active cases and their treatment under supervision is important. Poverty, poor hygiene, illiteracy, drug resistance, and poor compliance with medications are important reasons for the rising incidence of TB. Effective health education, access to treatment centers, and trained and motivated health care providers can go a long way in making national TB control programs a success. Acid fast staining of sputum is the best method if performed by experienced microbiologists, as it is reliable and economical. It can be said confidently that such kind of research will play a vital role in the fight against TB in Pakistan and especially in the district under study.

## REFERENCES

- Alli O, OD O, OO A (2011). Direct molecular detection of *Mycobacterium tuberculosis* complex from clinical samples: An adjunct to cultural method of laboratory diagnosis of tuberculosis. *North Am. J. Med. Sci.* 3(6): 281-288.
- Anand KM, Nag VL, Kant S, Kushwaha RAS, Kumar M, Mishra V, Rahman W, Tapan N, Dhole (2012). Evaluation of an immune chromatographic test for discrimination between *Mycobacterium tuberculosis* complex and non-tuberculous mycobacteria in clinical isolates from extra-pulmonary tuberculosis. *Ind. J. Med. Res.* 135: 901-906.
- Andersen P, Munk ME, Pollock JM, Doherty TM (2000). Specific immune- Alvi, A., S.F. Hussain, M.A. Shah, M. Khalida, M. Shamsudin. 1998. Prevalence of pulmonary tuberculosis on the roof of the world. *The Int. J. Tub. and Lung Dis.* 2(11):909-913.
- Broekmans JF, Migliori GB, Rieder HL (2002). European framework for tuberculosis control and elimination in countries with a low incidence: recommendations of the World Health Organization (WHO), International Union against Tuberculosis and Lung Disease and Royal Netherlands Tuberculosis Association Working Group.
- Bushra J, Qamruddin S, Sarwari AR, Hasan R (2008). An Assessment of Mantoux test in the Diagnosis of Tuberculosis in a BCG-Vaccinated, Tuberculosis-Endemic Area. *Infect. Dis. J. Pak.* 17 (1): 17-22.
- Dinesh D, Venkatachalam J, Singh Z, Joyce S, Anil J, Sathya GR (2013). Prevalence of Chest Symptomatics in a Rural Population of Kancheepuram District, South India. 3: 52-55
- Elena C, Cojocaru C, Saramet Bi, Diculencu D (2012). Identification of *Mycobacterium tuberculosis* complex using an immune chromatographic test. Running title: A useful test for *M. tuberculosis* identification. *Romanian Biotechnol. Letters.* 17(4): 7525.
- Inger B, Weldingh K, Lillebaek T, Follmann F, Andersen P (2004). Comparison of Tuberculin Skin Test and New Specific Blood Test in Tuberculosis Contacts. *Am. j. resp. crit. care medicine*, 170: 65-69.

- Javaid AK, Irfan M, Zaki A, Beg M, Hussain SF, Rizvi N (2006). Knowledge, Attitude and Misconceptions regarding Tuberculosis in Pakistani Patients. *J. Pak. Med. Assoc.* 56(5): 211-214.
- Nisar AR (2007). New drugs in resistant tuberculosis. *J. Pak. Med. Assoc.* 57(5): 252-6.No. 3
- Paige C, Bishai RW (2010). Penitentiary or penthouse condo: the tuberculous granuloma from the microbe's point of view. *Cell Microbiol.* 12(3):301-309.
- Rohit SS, Singla MN, Sharma PP (2001). Diagnosis of tuberculosis under rntcp: examination of two or three sputum specimens. *ind.j. tub.* 48: 13: 13-16.
- Shah SK, Sadiq H, Khalil M, Noor A, Rasheed G, Shah SM, Ahmad N (2003). Do private doctors follow national guidelines for managing pulmonary tuberculosis in Pakistan? *Eastern Mediterranean Health J.* 9(4): 776-88.
- Sun HJ, Golub JE, Jo J, Park S, Ohrr H, Samet JM (2009). Smoking and Risk of Tuberculosis Incidence, Mortality, and Recurrence in South Korean Men and Women. *Am. J. Epidemiol.* 170:1478-1485.
- WHO (2007). Tuberculosis Deaths Among Populations With High Hiv Prevalence.
- Young AK, Lee HYYH, Cho B, Han KS, Shim SY, Yim JJ (2005). Discrepancy Between the Tuberculin Skin Test and the Whole-Blood Interferony Assay for the Diagnosis of Latent Tuberculosis Infection in an Intermediate Tuberculosis-Burden Country. *JAMA;* 293(22):2756-2761.